

CRYPTOMAX™

Reg. No. / Nr L11248 Act No. 36 of 1947 / Wet Nr 36 van 1947

READ THE LABEL BEFORE USE
KEEP OUT OF REACH OF
CHILDREN AND ANIMALS

LIVE BIOLOGICAL CONTROL AGENT

STORE IN REFRIGERATOR
SHAKE WELL BEFORE USE

CRYPTOMAX™ is a suspension
concentrate biological agent for the
control of false codling moth larvae
on crops as listed.

CRYPTOMAX™ is 'n suspensie
konsentraat biologiese produk vir
die beheer van vals kodlingmot larwes
op gelyste gewasse.



WARNING

Hazard Statements
May cause an allergic skin reaction
Precautionary Statements
Avoid breathing spray
Wear protective clothing

ACTIVE INGREDIENT(S) / AKTIEWE BESTANDEEL(ELE):

Cryptophlebia leucotreta granulovirus (CrleGV-SA) at least
5 x 10¹⁰ occlusion bodies (OBs)/ml / ten minste 5 x 10¹⁰ okklusie partikels (OPs)/ml

Net mass/volume



Netto massa/volume

REGISTRATION HOLDER / REGISTRASIEHOUER

RIVER BIOSCIENCE (PTY) LTD | REG NO. 2003/022759/07 REG NR
PO BOX 20388, Humewood, 6013 | T: +27 41 373 1409
E: admin@riverbio.com | W: www.riverbioscience.co.za



Batch No.:

Date of Manufacture:



Expiry Date:

Shelf Life:

30 months from date of manufacture

: Lot Nr.

: Datum van Vervaardiging

: Vervaldatum

: Raklewe



2024.09.20 v2

WARNINGS

- May cause an allergic skin reaction.
- Store (in a refrigerator) away from food and animal feeds. CRYPTOMAX™ must be kept refrigerated (at approximately 4°C) until use. CRYPTOMAX™ must be used within 30 months of manufacture. While in use, the CRYPTOMAX™ container must be kept cool and in the shade. The container must not be exposed directly to sunlight.

DISCLAIMER

ALTHOUGH THIS REMEDY HAS BEEN EXTENSIVELY TESTED UNDER A LARGE VARIETY OF CONDITIONS, THE REGISTRATION HOLDER DOES NOT WARRANT THAT IT WILL BE EFFECTIVE UNDER ALL CONDITIONS, AS THE ACTION AND EFFECT THEREOF MIGHT BE AFFECTED BY FACTORS SUCH AS ABNORMAL CLIMATIC AND STORAGE CONDITIONS, QUALITY OF DILUTION WATER IN THE SPRAY TANK, COMPATIBILITY WITH OTHER SUBSTANCES NOT INDICATED ON THE LABEL AND THE OCCURRENCE OF RESISTANCE OF THE PEST AGAINST THE REMEDY CONCERNED AS WELL AS THE METHOD, TIME AND ACCURACY OF APPLICATION. FURTHERMORE, THE REGISTRATION HOLDER DOES NOT ACCEPT RESPONSIBILITY FOR DAMAGE TO CROPS, VEGETATION, THE ENVIRONMENT NOR HARM TO MAN OR ANIMAL. NEITHER DOES THE REGISTRATION HOLDER ACCEPT RESPONSIBILITY FOR LACK OF PERFORMANCE OF THE REMEDY CONCERNED, DUE TO FAILURE OF THE USER TO FOLLOW THE LABEL INSTRUCTIONS, NOR TO THE OCCURRENCE OF CONDITIONS WHICH COULD NOT HAVE BEEN FORESEEN IN TERMS OF THE REGISTRATION. CONSULT THE SUPPLIER IN THE EVENT OF ANY UNCERTAINTY.

PRECAUTIONS

- Avoid breathing spray. • Wear protective clothing.
- Contaminated work clothing should not be allowed out of the workplace.
- Do not eat, drink or smoke whilst mixing or applying, nor before washing hands and face. • Prevent contamination of food, feeds and drinking water.
- Prevent drift of spray mist onto other crops, grazing, rivers, dams, and areas not under treatment. • In case of spillage, absorb the liquid and scrub the area with 1-5% bleach solution. • Rinse empty container three times with a volume of water equal to 10% of the volume of the container. Add the rinsings to the contents of the spray tank. • Destroy empty container and do not use for any other purpose. • Dispose of in accordance with local regulations.

FIRST AID

- IF ON SKIN: Wash with plenty of water. Take off contaminated clothing and wash it before re-use. If skin irritation or rash occurs: Get medical help.
- Inhaled, move person to fresh air. • Rinse contaminated eyes with clean water for 15 minutes.

DIRECTIONS FOR USE - Use only as directed

CRYPTOMAX™ is a highly specific naturally occurring pathogen of the false codling moth. CRYPTOMAX™ does not harm bees, fish, birds, beneficial insects, or wildlife. CRYPTOMAX™ is fully compatible with an integrated pest management programme, as it is not harmful, nor disruptive to beneficial insects and other arthropods. Treated crops may be harvested shortly after application. There are no restrictions on applying CRYPTOMAX™ up to the time of harvest. Due to detrimentally high levels of UV-irradiation, the use of CRYPTOMAX™ is not recommended in the Clanwilliam and Pletberg magisterial districts of the Western Cape and the Northern Cape from December to March. Application should be conducted in these areas before and after the aforesaid period.

MIXING INSTRUCTIONS

Shake the container well before each use. For optimum performance, water pH should be between 5 and 8. Partially fill the spray tank with water, add the required amounts of CRYPTOMAX™ and adjuvants while agitating and top up the spray tank with water to the required volume.

APPLICATION INSTRUCTIONS

CRYPTOMAX™ should be applied immediately after mixing. Under no circumstances should it be left to stand overnight. CRYPTOMAX™ must be applied with an oscillating tower mist blower or with handheld spray guns. CRYPTOMAX™ must be applied as a full cover film spray. Do not concentrate the mixture in mist blowers. Due to the UV sensitivity of CRYPTOMAX™, it must not be applied during the heat of the day. Sprays must be applied during the late afternoon or at night.

RECOMMENDATIONS

PEST	CROP	RATE/100L WATER	REMARKS
False Codling Moth	Citrus	10ml + wetter*	Apply as a full cover film spray from when the pest first occurs. Sprays should be applied shortly after moth flight peaks. A final application 4 weeks before harvest is recommended. Apply during later afternoon or evening. In high pest pressure conditions, the application of molasses at 250ml per 100L water is recommended.

*Non-ionic organo-silicone or alkylated phenol-ethylene oxide wetters recommended at the registered rate such as, L6764 Break-Thru™, Polyether- polymethylsiloxane -copolymer, Evonik